

Accepted Manuscript

Advances in the Diagnosis and Management Of Asthma in Older Adults

Mazen Al-Alawi, MD, PhD Tidi Hassan, MD Sanjay H. Chotirmall, MD, PhD



PII: S0002-9343(13)01110-8

DOI: [10.1016/j.amjmed.2013.12.013](https://doi.org/10.1016/j.amjmed.2013.12.013)

Reference: AJM 12327

To appear in: *The American Journal of Medicine*

Received Date: 4 October 2013

Revised Date: 25 November 2013

Accepted Date: 2 December 2013

Please cite this article as: Al-Alawi M, Hassan T, Chotirmall SH, Advances in the Diagnosis and Management Of Asthma in Older Adults, *The American Journal of Medicine* (2014), doi: 10.1016/j.amjmed.2013.12.013.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**ADVANCES IN THE DIAGNOSIS AND MANAGEMENT
OF ASTHMA IN OLDER ADULTS**

Mazen Al-Alawi¹MD PhD, Tidi Hassan²MD and Sanjay H. Chotirmall³MD PhD

¹Department of Medicine, Our Lady of Lourdes Hospital, Navan, Republic of Ireland

²Department of Respiratory Medicine, Mater Misericordiae Hospital, Eccles Street, Dublin 7, Republic of Ireland

³Department of Medicine, St James's Hospital, James's Street, Dublin 8, Republic of Ireland

Corresponding Author:

Dr Sanjay H Chotirmall, Department of Medicine, St James's Hospital, James's Street, Dublin 8, Republic of Ireland

E-mail – schotirmall@rcsi.ie; Contact No. +353-87-9793833

Running Head: Asthma in older adults

Article Type: Review

Key Words: Asthma, Older adult, Elderly, Diagnosis, Treatment

Word Count: 3,916 (Abstract 174)

Conflicts of Interest: None of the authors have any conflicts of interest to disclose with respect to this manuscript

All authors has access to the data presented and a role in the preparation of the manuscript

Abstract

Global estimates on ageing predict an increased burden of asthma in the older population. Consequently, its recognition, diagnosis and management in clinical practice require optimization. This review aims to provide an update for clinicians highlighting advances in the understanding of the ageing process and immunosenescence together with their applicability to asthma from a diagnostic and therapeutic perspective. Ageing impacts airway responses, immune function and influences efficacy of emerging phenotype-specific therapies when applied to the elderly patient. Differentiating eosinophilic and neutrophilic disease accounts for atopic illness and distinguishes long-standing from late-onset asthma. Therapeutic challenges in drug delivery, treatment adherence and side effect profiles persist in the older patient while novel recording devices developed to aid detection of an adequate inhalation evaluates treatment effectiveness and compliance more accurately than previously attainable. Anti-cytokine therapies improve control of brittle asthma while bronchial thermoplasty is an option in refractory cases. Multi-dimensional intervention strategies prove best in the management of asthma in the older adult which remains a condition that is not rare but rarely diagnosed in this patient population.

Introduction

A World Health Organization (WHO) report on active ageing estimates that the proportion of individuals over the age of 65 is expected to more than double by 2050 ⁽¹⁾. The burden of respiratory disease is concurrently set to increase in the same period with an estimated 300 million people worldwide suffering from asthma, with 250,000 annual deaths attributed to the disease ⁽²⁾. Age-specific mortality associated with asthma in the older patient contrasts with the falling figures in younger age groups highlighting a basic need to improve asthma care and its management in the older population. Despite this, evidence from the TENOR study reports that hospitalization among older asthmatics remained lower than that of a younger population ⁽³⁾. This is in contrast to prospective data demonstrating that older patients with asthma were twice more likely to be hospitalized over a year-long follow-up period ⁽⁴⁾.

The National Institute on Aging (NIA) convened a workshop on asthma in the elderly that highlights differences in the pathophysiological mechanisms underpinning the disease in older patients that influence clinical course and outcomes ⁽⁵⁾. The aim of this current review is to provide an update for clinicians on the advances in understanding the biology and impact of the ageing process in regards to the diagnosis and treatment of asthma in the older adult. With major progress in the available and phenotype-specific therapies for asthma, its applicability to the older patient is described in the context of specific clinical challenges in this unique age group.

Long-standing versus Late-onset asthma

Asthma in the older adult is broadly divided into patients with long-standing disease present from childhood and, late-onset disease describing those developing symptoms following the sixth decade of life. The diagnosis of the latter is particularly challenging as its symptoms mimic alternative pathologies present in an older age-group such as chronic obstructive pulmonary disease or congestive cardiac failure (Tables 1 & 2). Airway inflammatory cell types determine the physiological responses observed and in the older asthmatic, eosinophilic inflammation is associated with airway hyper-responsiveness while neutrophils are an important determinant of airflow limitation at rest and during bronchoconstriction ⁽⁶⁾. Interestingly, chronic residential traffic pollution exposure is associated with eosinophilic but not neutrophilic inflammation in the older asthmatic ⁽⁷⁾. These findings strengthen the relationship between airway pathophysiology and clinical phenotypic differences observed in the older patient with asthma.

The biology of ageing and its effect on the diagnosis of asthma

Ageing is the natural process of physiological change occurring within organ systems decreasing their functional capacity. This in turn increases risk of disease. Whilst all individuals undergo this process, vast heterogeneity exists making the impact of a particular disease having the potential to manifest differently depending on molecular, epigenetic and individual factors. Environmental insults when combined with reduced capacity for DNA repair with ageing increases the fragility of the lung to regenerate. Like other organs, the lung continually loses capacity over time resulting in compromised pulmonary function. We have previously highlighted the inherent difficulties in performing and interpreting pulmonary function testing in

older patients because of the effects of age-related changes in lung function on respiratory physiology⁽⁸⁾. Aging leads to an obstructive defect on pulmonary function testing that may be challenging in distinguishing from a superimposed active disease process such as asthma. Therefore a combination of clinical history, physical examination and diagnostic testing are critical to achieve an appropriate diagnosis that in turn directs treatment.

Clinical Assessment

Asthma is a heterogeneous disease entity with increasing emergence of varying clinical phenotypes. One major differentiation of relevance to the older patient is that between longstanding and late-onset asthma⁽⁸⁾. The clinical history remains pivotal to aid diagnosis. For example, allergic nasal symptoms diminish with age hence making a history of allergy less useful in this age group. A definitive history on environmental exposures and irritants such as cigarette smoking, household aerosols, paints, perfumes and inhaled metabisulfites found in beer and food preservatives is invaluable. Another critical aspect includes medication history such as aspirin, non-steroidal anti-inflammatories, angiotensin converting enzyme inhibitors, beta-blockers and hormonal administration that could potentially induce bronchoconstriction. Differentiating asthma from other co-morbidities is by far the greatest clinical challenge as cardiac failure, chronic aspiration, upper airway obstruction, reflux disease and chronic obstructive pulmonary disease (COPD) all mimic asthma (Tables 1 & 2).

Asthma in the older adult displays many of the hallmarks of COPD such as onset of symptoms later in life, partial reversibility on pulmonary function testing and association with neutrophilic inflammation⁽⁹⁾. Co-existing COPD with an asthmatic

phenotype is encountered and poses an additional diagnostic challenge. Other factors accounting for a delayed diagnosis include poorer perception of dyspnea in the elderly and the psychosocial impact of aging. This results in aberrant reporting of symptoms exacerbated by the co-existence of depression, cognitive impairment, social isolation, and denial.

Physiological Assessment

Demonstrating physiological impairment assists the diagnosis of asthma. However, significant variability exists in the ability to detect airway obstruction in the older asthmatic. Initial spirometry may be normal and follow-up testing necessary for detecting airway obstruction ⁽⁵⁾. The clinical utility of pulmonary function tests are user dependent: the inability to follow instructions due to poor coordination or cognitive impairment renders performance and subsequently interpretation of results challenging in this cohort ⁽⁸⁾. One alternative is the measurement of respiratory impedance. This simple technique requiring minimal patient cooperation has been used successfully in young children and older adults for both diagnosis and therapeutic monitoring of respiratory symptoms associated with airway obstruction ⁽¹⁰⁾. Assessment of institutionalized patients with cognitive impairment highlighted the superiority of respiratory impedance over spirometry in this patient group ⁽¹¹⁾. Home peak flow monitoring has been suggested as both a diagnostic and monitoring tool for asthma, however compliance is often challenging in the older patient ⁽¹²⁾. Although significant early morning dips aid diagnosis, high variability remains a poor predictor of asthma and hence it's limited role in late-onset disease ⁽¹³⁾.

Ageing and the Airway response

The ageing process impacts upon airway responses utilized in the diagnosis of asthma. Ageing however does not alter the degree of response to inhaled bronchodilator drugs. The use of combined albuterol in conjunction with ipratropium bromide for diagnostic testing in the older adult often produces a more effective degree of bronchodilation. It is important to note that the time to achieve peak effect in bronchodilation is 30 minutes in contrast to 5-10 minutes when using albuterol as a single agent bronchodilator^(14, 15). The changes in calibre of small airways in response to bronchodilation should not be used in the elderly asthmatic as it often increases in response to the reduction of air trapping within the lungs⁽¹⁶⁾.

Gronke *et al.* (2002) identified that airway hyper-responsiveness was evident in individuals with a short (≤ 16 years) but not with long (≥ 16 years) duration of asthma⁽¹⁷⁾. This suggests that methacholine challenge testing, an accurate diagnostic tool for asthma in symptomatic patients with normal pulmonary function tests, may be of greater use in late onset versus long-standing asthma. Furthermore, the prevalence of hyper-responsiveness is greater in older adults despite correcting for atopy, degree of airway obstruction and smoking history suggesting that a lower provocative challenge dose ($<4\text{mg/mL}$) may be more appropriate to define airway hyper-responsiveness in the older patient than the conventional $<8\text{mg/mL}$ ⁽¹⁸⁾.

Therapeutics and its associated challenges in the older adult

Effective asthma management in the older patient relies on similar principles applicable to all ages. Key features include education; monitoring and effective control of environmental factors in addition to pharmacological therapy. The National Asthma Education and Prevention Program recommend a stepwise approach to therapy. A number of therapies are generally not recommended for use in the elderly asthmatic. Zileuton, an inhibitor of 5-lipoxygenase necessitates a regular monitoring of liver function whilst theophylline's wide range of drug interactions affect serum concentration and subsequent toxicity precludes its routine use in the elderly patient.

Asthma in the older adult assumes a unique phenotype, and accordingly we propose an algorithm adapted from the stepwise Global Initiative for Asthma (GINA) guidelines to aid clinical decision making when managing asthma in an older individual ⁽¹⁹⁾ (Figure 1).

In addition, there is emerging evidence for the role of long acting anticholinergic inhalers in the management of asthma. Bateman *et al* report that tiotropium was not inferior to salmeterol in maintaining an improved lung function in asthmatics ⁽²⁰⁾. Peters *et al* explored the role of clinical parameters predicting a response to tiotropium therapy ⁽²¹⁾. They identified that an acute response to short-acting bronchodilators predicted a positive clinical response to tiotropium. Despite this, a meta-analysis carried out by Tian *et al* highlighted that the duration of most trials was too short to allow for evaluation of long-term efficacy and safety of tiotropium ⁽²²⁾. It is also important to recognize that current trials investigating the use of tiotropium have evaluated its role as additional therapy to current management guidelines rather than a first-choice stand-alone therapeutic option. Future longer-term studies are expected to address this. It is likely however that tiotropium therapy

will provide clinicians with a future option in controlling symptoms in the elderly asthmatic by concomitant treatment of age-related COPD in a convenient once daily dosing regimen.

The role of long-term macrolide therapy to improve asthma control has been explored in a number of trials. A meta-analysis of four randomised controlled trials identified improvement in symptoms, quality of life and airway hyper-reactivity, but highlighted the limited statistical power to detect significant differences in lung function⁽²³⁾.

If considered for use in the elderly asthmatic, it is critical to highlight a number of complex drug interactions occurring between macrolide therapy and a number of pharmacological agents used for symptom maintenance including anti-histamines and theophyllines (Table 3). Future longer-term trials focusing on both the safety and efficacy of macrolides on exacerbation rates may highlight these agents as a useful adjunct in the management of late-onset asthma.

β -blockade therapy may exacerbate symptoms of asthma when administered topically for glaucoma or orally for acute coronary syndromes. Initiating β -blocker therapy in the older asthmatic must be performed under close supervision as severe life-threatening symptoms may be triggered by the initial dose^(24, 25). There is however evidence suggesting that escalating β -blocker therapy in asthmatics is well tolerated⁽²⁶⁾. The risk-benefit ratio of β -blocker maintenance therapy in asthmatics has a limited evidence base. This is further complicated by potential beneficial effects on airway inflammation and hyper-responsiveness in some patients with asthma⁽²⁷⁻²⁹⁾. Clinical benefits of selective β -blocker therapy in patients with coronary disease necessitates a careful assessment by the prescribing clinician with regards the presence of concomitant poorly controlled asthma or the potential risks of poor

symptom control following initiation. Holding β -blockade may be appropriate in the setting of hospital admission for decompensated asthmatic symptoms in the older adult but the relative risk-benefit ratio must be considered in terms of other co-morbid illness.

Drug delivery and adherence

Despite the fundamental role of inhaled bronchodilators and corticosteroids many elderly patients remain undertreated⁽³⁰⁾. Early work estimates that half of all patients are unable to use the inhalers prescribed and despite increased awareness only marginal improvements have occurred^(31, 32). Availability of preconstituted combination therapies and spacer devices has increased clinical efficacy and treatment adherence. Limitations to the latter do remain owing to cumbersome sizes and difficulty with maintenance of a sterile spacer device key to eliminate unfavorable electrostatic charges⁽³³⁾. In an older patient, maintaining a clean spacer is particularly challenging as contamination due to colonization with *Pseudomonas aeruginosa*, methicillin-resistant *Staphylococcus aureus*, *Burkholderia*, and *Stenotrophomonas maltophilia* has been reported⁽³⁴⁾.

Major technological advances have occurred in the field of assessing inhaler compliance and adherence to prescribed treatment. Such methods have the capacity to revolutionize treatment compliance and effectiveness in the older population. Many older patients are unable to achieve a peak inspiratory flow rate necessary to fully extract medication from a dry-powdered inhaler. Novel recording devices have been developed to aid identification and detection of an adequate inhalation utilizing acoustic signals⁽³⁵⁾.

Treatment effectiveness

Effectiveness of therapy in the elderly population has only been investigated on a trial basis by two current therapies. The ACCEPT trial explored the effect of age on the response to zafirlukast. It failed to demonstrate a significant change in spirometric decline and rescue inhaler use with zafirlukast however highlighted clinically significant differences in asthma symptoms⁽³⁶⁾. Omalizumab therapy in an elderly veteran population demonstrated significant positive clinical responses where moderate numbers of patients on long-term corticosteroids were able to discontinue treatment⁽³⁷⁾.

A limited pool of available evidence highlights a growing need to evaluate therapies and treatment algorithms for older asthmatics in specific trials fit for this purpose. A major factor contributing to the lack of available evidence in the elderly asthmatic population remains the screening techniques employed by large multi-centre work. Older patients are frequently excluded from such investigational work due to burden of comorbidity and the need to simplify the measured outcomes of clinical trials⁽³⁸⁾. The screening criteria employed in clinical trials for asthma inherently exclude the elderly asthmatic as phenotypic changes in the degree of airway hyper-responsiveness, reductions in eosinophil counts and function, and change in fractional exhaled nitric oxide (FeNO) responses render the elderly individual ineligible for inclusion. Dupilumab, an IL-4 monoclonal antibody has been recently trialled for persistent asthma with elevated eosinophil counts and within the study a number of Th2 biomarkers were analysed at various time points⁽³⁹⁾. Any patient over 65 was excluded from the trial as surrogate markers used to monitor response to treatment are influenced by ageing. Serum biomarkers including thymus

activation-regulated chemokine (TARC), immunoglobulin-E (IgE), FeNO and blood/sputum eosinophils are impacted upon by immunosenescence. Future research initiatives should address such inherent problems in patient selection and outcome measures during asthmatic trials. Differences between older and younger asthmatics need to be sought in the appropriate clinical context, limited at present because of a lack of enrolment within asthma trials.

Pathophysiological impact of ageing and effects on emerging asthma therapies

Atopic Asthma

Aging is associated with a decline in the prevalence of atopic symptoms, IgE levels and positive skin allergen tests. Healthy older adults have at least a single positive allergen skin test in contrast to the older patient with asthma that develops multiple sensitisations to common indoor allergens such as cats, dogs, mites and cockroaches^(40, 41). Older patients that may have become sensitised to cockroaches tend to develop more severe asthma and experience a steeper decline in lung function^(42, 43). Places of residence are known to influence the frequency of acute respiratory infections from 1-2% per year to 6-11% per year at day-care and long-term centres⁽⁴⁴⁾. Clinical utility of controlling triggers in the elderly with measures to control aeroallergen exposure to known agents should be initiated. Some of the common provocative agents include cigarette smoke, paints, varnish, and cleaning aerosols.

The eosinophil remains the cellular conductor of allergic asthma and eosinophil counts remain comparable between older and younger asthmatics. Degree of degranulation is however significantly reduced in the elderly patient. A critical

limitation of published work to date remains the use of peripheral blood rather than airway eosinophils. The two populations exhibit crucial differences in cell surface markers and activation states that remain unaddressed by use of sole peripheral sampling⁽⁴⁵⁾.

Furthermore, an inverse relationship between ageing and IgE production exists suggestive of a natural desensitization process. Current evidence suggests that development of asthma in the older patient is not linked to total IgE but rather to sensitization and the development of allergen-specific IgEs following a prolonged environmental exposure to allergens. This remains an important consideration when evaluating the use of anti-IgE strategies in elderly asthmatics. A potential false elevation in total serum IgE must be recognized confounded by factors including sex, smoking status, concurrent infections and environmental sensitisation^(6, 46). During both the diagnosis and more importantly escalation of therapy to include the anti-IgE agent Omalizumab, such features must be taken into account. Omalizumab therapy should however always be considered in the elderly asthmatic as a safe escalation option but confounding total IgE levels may complicate its initiation and possibly effectiveness. Importantly however, Omalizumab has been reported to reduce exacerbations, improve symptoms and limit the need for rescue medication in asthmatics. Its global effectiveness reported by both clinicians and patients extends to elderly patients with moderate-severe allergic asthma and remains, in the right circumstances a viable therapeutic option⁽³⁷⁾.

Non-atopic asthma

One of the key advances in the recent era of asthma care has been novel approaches to testing for airway inflammation. This allows differentiation between eosinophilic, neutrophilic or lymphocytic based disease differentiating atopic from non-atopic asthma but additionally from COPD.

Nitric oxide is generated by a variety of inflammatory cells that include polymorphonuclear leukocytes, mononuclear cells, and eosinophils. Identification of nitric oxide in exhaled breath may predict asthma exacerbations⁽⁴⁷⁾. Hardaker *et al* (2011) highlighted that although exhaled nitric oxide is a strong predictor of asthma severity in the younger patient, this correlation disappears rapidly with age⁽⁴⁸⁾. Clinical utility of exhaled nitric oxide is currently limited by absence of evidence to determine if it is causative of airway dysfunction or a marker of a normal homeostatic mechanism⁽⁴⁹⁻⁵²⁾. The development of non-invasive bed-side testing for airway inflammation such as analysis of breath condensate cytokines and induced sputum will allow differentiation between eosinophilic, neutrophilic, and lymphocytic bronchial inflammation⁽⁵³⁾. These may be of greater clinical value than exhaled nitric oxide in the older adult allowing a directed and more personalized therapeutic approach.

Most reflective of late-onset disease, non-atopic asthma is often triggered by viral upper respiratory tract infections which leads to exacerbations. Detecting exacerbations early requires clinical acumen in this population because of relatively poor distinguishing features from COPD and a lack of sensitive viral diagnostic testing⁽⁵⁾. What remains intriguing in the older adult is whether viral exacerbations occur secondary to persistent airway inflammation or age-related immune-

inflammatory changes termed immunosenescence. This active and highly regulated process involves bidirectional crosstalk between hematopoietic cells and thymic epithelia⁽⁵⁴⁾. Whilst established evidence point toward a Th2-cytokine bias in early life, studies in older adults indicate a key role for Th1 responses providing further evidence for age associated changes in the airway inflammatory response⁽⁵⁵⁾. Furthermore, murine models of asthma in the context of ageing have demonstrated diminished B-cell populations and a transition from naïve to antigen-expressing B-cells⁽⁵⁶⁾. Parallel reductions in antibody production may be responsible for the enhanced antigen persistence and specificity observed in the elderly. Thymic involution evokes shifts in the T-cell population, alterations in antigen B-cell processing, eosinophil function, and reduction in phagocytic capabilities, all of which contribute to a unique immunological milieu in the older adult with asthma.

Anti-cytokine therapies

In the context of immunosenescence, promising avenues of therapeutics targeting inflammatory cascades continue to emerge. Asthmatic airways contain CD4+ T-cells that produce IL-4 and -13 which are both therapeutic targets that have been investigated. The range of antigenic activators associated with the CD4 pathway is continually being expanded and this group of therapeutics may yet provide novel avenue streams to treat the elderly asthmatic. Serum obtained from older asthmatics demonstrates increased concentration and persistence of certain T-cell subsets highlighting the key role of prolonged activation and survival of these cell types in the pathophysiology of late-onset asthma. This expanding arsenal of anti-cytokine therapies utilized in poorly controlled asthma is summarized (Table 4) and despite a lack of data in older patients provides the most promising newer therapeutic options.

It must however be noted that the emergence of anti-cytokine therapy does not come without risk as IL-4, -5, and eosinophils have important innate immunological roles. Development of serious infections and neoplasia both at higher risk in an elderly population halted the development of golimumab (anti-TNF- α antibody) and highlighted the key protective effects of IL-17A and -17F in immunologic surveillance⁽⁵⁷⁾. The true risk-benefit effects of these emerging agents applicable to the older asthmatic are yet to be fully elucidated.

Bronchial thermoplasty

Bronchial thermoplasty is a treatment option for the poorly controlled asthmatic refractory to other therapies. The technique involves endoscopic application of thermal radiofrequency energy to ablate underlying smooth muscle altering airway structure. It has been shown that structural changes in the older fatal asthmatic share overlapping features with those of younger fatalities⁽⁵⁸⁾. This suggests that bronchial thermoplasty may be of benefit in a selected subset of older patients with asthma. However, the safety and feasibility of the procedure remains to be determined in an elderly population despite bronchoscopy being reported as a relatively safe and well tolerated procedure in octogenarians⁽⁵⁹⁾.

Multi-dimensional interventional strategies and pulmonary rehabilitation

Use of a multidimensional assessment and intervention strategy has been highlighted for effective management of asthma in the older adult⁽⁶⁰⁾. This approach validated by systemic reviews and randomized controlled trials identify several dimensions of care including pharmacotherapy, individual rehabilitation and social interventions. By addressing specific age-related issues such as comorbidities and acute care

complications, the strategy contributes to an improved quality of life and cost effectiveness if domains are targeted appropriately. Owing to its success in managing asthma, it has been adapted into the management of COPD affecting the older adult ⁽⁶¹⁾. Additionally, the role of pulmonary rehabilitation programs in the management of asthma remains to be fully appreciated. Cox *et al* initially highlighted a role for rehabilitation in a small population of patients with asthma illustrating improvements in symptom control and quality of life ⁽⁶²⁾. Laurino *et al* highlighted that breathing retraining improved symptoms of asthma related to anxiety and subsequently improved quality of life ⁽⁶³⁾. Despite these small studies, the longer-term beneficial effects of a pulmonary rehabilitation program particularly on an elderly asthmatic population remain to be determined.

Conclusion

Despite advances in both diagnostics and therapeutics, asthma remains an under-recognized health issue in the elderly population. This diagnosis is not rare but it is rarely diagnosed as almost ten percent of the adult population over 65 years of age are afflicted. Emerging evidence suggests that asthma in the older adult is phenotypically distinct from that seen in younger patients particularly when physiological changes associated with the ageing process are taken into account. In addition to an increased number of elderly patients being enrolled into asthma trials, dedicated and well-funded clinical studies of this population are required to evaluate future diagnostic and treatment approaches. Research in the field needs to focus on both human and animal model systems to investigate the impact of the ageing process on the immunologic pathways underpinning asthma in the ageing lung. Epigenetic, environmental and microbiological triggers need to be considered in the clinical

setting while translational investigation of novel therapeutic targets must be pursued. Our current knowledge base is largely from specimens obtained from young adults however the role of allergic asthma diminishes with age hence a different phenotype presents in the elderly with normal eosinophils and elevated sputum neutrophils. Neutrophilic asthma is associated with innate immune pathways in contrast to Th2 mediated allergic asthma hence research on alternative signalling cascades involving NKT/Th17 lymphocyte subtypes must be pursued. Future work should not be limited to therapeutic targets but also identify non-invasive biomarkers of disease severity and progression. Clinical trials in this challenging population need to account for the unique phenotypic makeup of the elderly asthmatic and for the immunological and physiological changes associated with the natural process of ageing that affect its diagnosis and management.

References

1. Kalache A, Keller I. The WHO perspective on active ageing. *Promot Educ.* 1999;6(4):20-3, 44, 54.
2. Bousquet J, Dahl R, Khaltaev N. Global Alliance against Chronic Respiratory Diseases. *Eur Respir J.* 2007;29(2):233-9.
3. Chipps BE, Zeiger RS, Borish L, et al. Key findings and clinical implications from The Epidemiology and Natural History of Asthma: Outcomes and Treatment Regimens (TENOR) study. *J Allergy Clin Immunol.* 2012;130(2):332-42 e10.
4. Diette GB, Krishnan JA, Dominici F, et al. Asthma in older patients: factors associated with hospitalization. *Arch Intern Med.* 2002;162(10):1123-32.
5. Hanania NA, King MJ, Braman SS, et al. Asthma in the elderly: Current understanding and future research needs--a report of a National Institute on Aging (NIA) workshop. *J Allergy Clin Immunol.* 2011;128(3 Suppl):S4-24.
6. Porsbjerg CM, Gibson PG, Pretto JJ, et al. Relationship between airway pathophysiology and airway inflammation in older asthmatics. *Respirology.* 2013.
7. Epstein TG, Kesavalu B, Bernstein CK, et al. Chronic traffic pollution exposure is associated with eosinophilic, but not neutrophilic inflammation in older adult asthmatics. *J Asthma.* 2013;50(9):983-9.
8. Chotirmall SH, Watts M, Branagan P, et al. Diagnosis and management of asthma in older adults. *J Am Geriatr Soc.* 2009;57(5):901-9.
9. Nyenhuis SM, Schwantes EA, Evans MD, Mathur SK. Airway neutrophil inflammatory phenotype in older subjects with asthma. *J Allergy Clin Immunol.* 2010;125(5):1163-5.

10. Frei J, Jutla J, Kramer G, et al. Impulse oscillometry: reference values in children 100 to 150 cm in height and 3 to 10 years of age. *Chest*. 2005;128(3):1266-73.
11. Carvalhaes-Neto N, Lorino H, Gallinari C, et al. Cognitive function and assessment of lung function in the elderly. *Am J Respir Crit Care Med*. 1995;152(5 Pt 1):1611-5.
12. Dow L, Fowler L, Phelps L, et al. Prevalence of untreated asthma in a population sample of 6000 older adults in Bristol, UK. *Thorax*. 2001;56(6):472-6.
13. Enright PL. The diagnosis and management of asthma is much tougher in older patients. *Curr Opin Allergy Clin Immunol*. 2002;2(3):175-81.
14. Kradjan WA, Driesner NK, Abuan TH, et al. Effect of age on bronchodilator response. *Chest*. 1992;101(6):1545-51.
15. Hardie JA, Buist AS, Vollmer WM, et al. Risk of over-diagnosis of COPD in asymptomatic elderly never-smokers. *Eur Respir J*. 2002;20(5):1117-22.
16. Lung function testing: selection of reference values and interpretative strategies. American Thoracic Society. *Am Rev Respir Dis*. 1991;144(5):1202-18.
17. Gronke L, Kannies F, Holz O, et al. The relationship between airway hyper-responsiveness, markers of inflammation and lung function depends on the duration of the asthmatic disease. *Clin Exp Allergy*. 2002;32(1):57-63.
18. Cuttitta G, Cibella F, Bellia V, et al. Changes in FVC during methacholine-induced bronchoconstriction in elderly patients with asthma: bronchial hyperresponsiveness and aging. *Chest*. 2001;119(6):1685-90.
19. Bateman ED, Hurd SS, Barnes PJ, et al. Global strategy for asthma management and prevention: GINA executive summary. *Eur Respir J*. 2008;31(1):143-78.

20. Bateman ED, Kornmann O, Schmidt P, et al. Tiotropium is noninferior to salmeterol in maintaining improved lung function in B16-Arg/Arg patients with asthma. *J Allergy Clin Immunol*. 2011;128(2):315-22.
21. Peters SP, Bleecker ER, Kunselman SJ, et al. Predictors of response to tiotropium versus salmeterol in asthmatic adults. *J Allergy Clin Immunol*. 2013;132(5):1068-74 e1.
22. Tian JW, Chen JW, Chen R, Chen X. Tiotropium versus placebo for inadequately controlled asthma: a meta-analysis. *Respir Care*. 2013.
23. Reiter J, Demirel N, Mendy A, et al. Macrolides for the long-term management of asthma - a meta-analysis of randomized clinical trials. *Allergy*. 2013;68(8):1040-9.
24. Raine JM, Palazzo MG, Kerr JH, Sleight P. Near-fatal bronchospasm after oral nadolol in a young asthmatic and response to ventilation with halothane. *Br Med J (Clin Res Ed)*. 1981;282(6263):548-9.
25. Spitz DJ. An unusual death in an asthmatic patient. *Am J Forensic Med Pathol*. 2003;24(3):271-2.
26. Arboe B, Ulrik CS. Beta-blockers: friend or foe in asthma? *International journal of general medicine*. 2013;6:549-55.
27. Hanania NA, Singh S, El-Wali R, et al. The safety and effects of the beta-blocker, nadolol, in mild asthma: an open-label pilot study. *Pulm Pharmacol Ther*. 2008;21(1):134-41.
28. Hanania NA, Mannava B, Franklin AE, et al. Response to salbutamol in patients with mild asthma treated with nadolol. *Eur Respir J*. 2010;36(4):963-5.

29. Callaerts-Vegh Z, Evans KL, Dudekula N, et al. Effects of acute and chronic administration of beta-adrenoceptor ligands on airway function in a murine model of asthma. *Proc Natl Acad Sci U S A*. 2004;101(14):4948-53.
30. Sin DD, Tu JV. Underuse of inhaled steroid therapy in elderly patients with asthma. *Chest*. 2001;119(3):720-5.
31. Crompton GK. Problems patients have using pressurized aerosol inhalers. *Eur J Respir Dis Suppl*. 1982;119:101-4.
32. Crompton GK, Barnes PJ, Broeders M, et al. The need to improve inhalation technique in Europe: a report from the Aerosol Drug Management Improvement Team. *Respir Med*. 2006;100(9):1479-94.
33. Lipworth BJ, Lee DK, Anhoj J, Bisgaard H. Effect of plastic spacer handling on salbutamol lung deposition in asthmatic children. *Br J Clin Pharmacol*. 2002;54(5):544-7.
34. Berlinski A, Waldrep JC. Nebulized drug admixtures: effect on aerosol characteristics and albuterol output. *J Aerosol Med*. 2006;19(4):484-90.
35. Holmes MS, Seheult JN, Geraghty C, et al. A method of estimating inspiratory flow rate and volume from an inhaler using acoustic measurements. *Physiological measurement*. 2013;34(8):903-14.
36. Korenblat PE, Kemp JP, Scherger JE, et al. Effect of age on response to zafirlukast in patients with asthma in the Accolate Clinical Experience and Pharmacoeconomics Trial (ACCEPT). *Ann Allergy Asthma Immunol*. 2000;84(2):217-25.
37. Maykut RJ, Kianifard F, Geba GP. Response of older patients with IgE-mediated asthma to omalizumab: a pooled analysis. *J Asthma*. 2008;45(3):173-81.

38. Mathur SK. Allergy and asthma in the elderly. *Semin Respir Crit Care Med*. 2010;31(5):587-95.
39. Wenzel S, Ford L, Pearlman D, et al. Dupilumab in persistent asthma with elevated eosinophil levels. *N Engl J Med*. 2013;368(26):2455-66.
40. Huss K, Naumann PL, Mason PJ, et al. Asthma severity, atopic status, allergen exposure and quality of life in elderly persons. *Ann Allergy Asthma Immunol*. 2001;86(5):524-30.
41. Barbee RA, Kaltenborn W, Lebowitz MD, Burrows B. Longitudinal changes in allergen skin test reactivity in a community population sample. *J Allergy Clin Immunol*. 1987;79(1):16-24.
42. Thorn J, Brisman J, Toren K. Adult-onset asthma is associated with self-reported mold or environmental tobacco smoke exposures in the home. *Allergy*. 2001;56(4):287-92.
43. Jaakkola MS, Nordman H, Piipari R, et al. Indoor dampness and molds and development of adult-onset asthma: a population-based incident case-control study. *Environ Health Perspect*. 2002;110(5):543-7.
44. Falsey AR, McCann RM, Hall WJ, et al. Acute respiratory tract infection in daycare centers for older persons. *J Am Geriatr Soc*. 1995;43(1):30-6.
45. Sedgwick JB, Calhoun WJ, Vrtis RF, et al. Comparison of airway and blood eosinophil function after in vivo antigen challenge. *Journal of immunology*. 1992;149(11):3710-8.
46. Viswanathan RK, Mathur SK. Role of allergen sensitization in older adults. *Current allergy and asthma reports*. 2011;11(5):427-33.

47. Taddei S, Galetta F, Viridis A, et al. Physical activity prevents age-related impairment in nitric oxide availability in elderly athletes. *Circulation*. 2000;101(25):2896-901.
48. Hardaker KM, Downie SR, Kermode JA, et al. Predictors of airway hyperresponsiveness differ between old and young patients with asthma. *Chest*. 2011;139(6):1395-401.
49. Baraldi E, Carra S, Dario C, et al. Effect of natural grass pollen exposure on exhaled nitric oxide in asthmatic children. *Am J Respir Crit Care Med*. 1999;159(1):262-6.
50. de Gouw HW, Hendriks J, Woltman AM, et al. Exhaled nitric oxide (NO) is reduced shortly after bronchoconstriction to direct and indirect stimuli in asthma. *Am J Respir Crit Care Med*. 1998;158(1):315-9.
51. Olin AC, Rosengren A, Thelle DS, et al. Height, age, and atopy are associated with fraction of exhaled nitric oxide in a large adult general population sample. *Chest*. 2006;130(5):1319-25.
52. Smith AD, Cowan JO, Brassett KP, et al. Use of exhaled nitric oxide measurements to guide treatment in chronic asthma. *N Engl J Med*. 2005;352(21):2163-73.
53. Sutherland ER, Martin RJ. Airway inflammation in chronic obstructive pulmonary disease: comparisons with asthma. *J Allergy Clin Immunol*. 2003;112(5):819-27; quiz 28.
54. Gill J, Malin M, Sutherland J, et al. Thymic generation and regeneration. *Immunol Rev*. 2003;195:28-50.

55. Sakuishi K, Oki S, Araki M, et al. Invariant NKT cells biased for IL-5 production act as crucial regulators of inflammation. *Journal of immunology*. 2007;179(6):3452-62.
56. Johnson SA, Rozzo SJ, Cambier JC. Aging-dependent exclusion of antigen-inexperienced cells from the peripheral B cell repertoire. *Journal of immunology*. 2002;168(10):5014-23.
57. Wenzel SE, Barnes PJ, Bleecker ER, et al. A randomized, double-blind, placebo-controlled study of tumor necrosis factor-alpha blockade in severe persistent asthma. *Am J Respir Crit Care Med*. 2009;179(7):549-58.
58. Senhorini A, Ferreira DS, Shiang C, et al. Airway dimensions in fatal asthma and fatal COPD: overlap in older patients. *COPD*. 2013;10(3):348-56.
59. Chotirmall SH, Watts M, Moore A, et al. Dispelling myths regarding the safety of 'bronchoscopy in octogenarians'. *Age Ageing*. 2009;38(6):764-5.
60. Gibson PG, McDonald VM, Marks GB. Asthma in older adults. *Lancet*. 2010;376(9743):803-13.
61. McDonald VM, Simpson JL, Higgins I, Gibson PG. Multidimensional assessment of older people with asthma and COPD: clinical management and health status. *Age Ageing*. 2011;40(1):42-9.
62. Cox NJ, Hendricks JC, Binkhorst RA, van Herwaarden CL. A pulmonary rehabilitation program for patients with asthma and mild chronic obstructive pulmonary diseases (COPD). *Lung*. 1993;171(4):235-44.
63. Laurino RA, Barnabe V, Saraiva-Romanholo BM, et al. Respiratory rehabilitation: a physiotherapy approach to the control of asthma symptoms and anxiety. *Clinics*. 2012;67(11):1291-7.

FIGURE LEGEND

Figure 1: Proposed stepwise approach to management of asthma in the older adult. (Adapted from GINA guidelines) SABA: short acting β -agonist; PRN: as required; ICS: inhaled corticosteroid; LTRA: leukotriene receptor antagonist; LABA: long acting β -agonist.

CLINICAL SIGNIFICANCE

- Asthma in older adults remains a condition not rare but rarely diagnosed
- Recognition, diagnosis and management in clinical practice require optimization
- Major challenges exist in performing and interpreting diagnostic testing and overcoming challenging therapeutic issues unique to this age group
- This review provides an update on the advances in the biology and impact of the ageing process with regard to the diagnosis and treatment of asthma in the older adult

	COPD	Long-standing Asthma (LSA)	Late-onset Asthma (LOA)
Spirometry	Obstructive	Obstructive	Normal or obstructive
*Reversibility	None or Minimal	Significant (12% change in FEV ₁ or 200 mL)	Significant (12% change in FEV ₁ or 200 mL)
Diffusing Capacity	Decreased	Normal	Normal
Hyperinflation	Increased	Normal or Increased during attacks	Normal or Increased during attacks

Table 1: Summary of physiological comparisons between asthma and chronic obstructive pulmonary disease (COPD). *Reversibility is measured by change in forced expiratory volume in 1-second (FEV₁) to bronchodilator therapy (salbutamol).

	Younger Asthmatic	Older Asthmatic
Allergic symptoms	Present	Likely absent
Airway responsiveness	Significant	Significant
Reversibility	Short acting β 2-agonist	Short acting β 2-agonist +/- Anti-cholinergic
Time to achieve peak bronchodilation	5-10 minutes	Up to 30 minutes
IgE	Normal or elevated	Total IgE (likely normal) and Allergen specific IgE (may be elevated)
Eosinophil Counts	Normal or elevated	Likely normal
Airway inflammation	Eosinophilic	Neutrophilic
Co-morbidities	Absent	COPD and/or CCF (most common)

Table 2: Summary of the key features to distinguish younger and older asthmatics.

IgE: Immunoglobulin-E; COPD: Chronic Obstructive Pulmonary Disease; CCF: Congestive Cardiac Failure.

Class	Drug	Mechanism	Effect
Antihistamines	Terfenadine	CYP 3A4	QT prolongation
	Loratadine	CYP 3A4 CYP 2D6	QT prolongation
Theophyllines	Aminophylline	CYP 1A2	Decreased theophylline
		CYP 3A4	excretion by 25%

Table 3: Macrolide therapy drug interactions with antihistamines and theophyllines.

CYP: Cytochrome families and relevant isoforms.

Target cytokine	Function	Therapeutic agent(s)
IL-2	Activating factor: Th1 & Th2 cells	Daclizumab
IL-5	Eosinophil growth, maturation and activation	Mepolizumab Reslizumab Benralizumab
IL-9	Mast cell proliferation and mucus hyperplasia	MEDI-528
IL-4	Th2 differentiation, expansion, and isotype switching Eosinophil recruitment, Mast cell development	Pitrakinra Dupilumab
IL-13	Bronchial fibroblast and airway smooth muscle proliferation, Eosinophil and Basophil recruitment	Lebrikizumab Anrukinzumab Tralokinumab
IL-17	Airway recruitment of Neutrophils, Lymphocytes and Eosinophils	Secukinumab
Anti-GM-CSF	Eosinophil differentiation and survival	MT203

Table 4: Emerging anti-cytokine therapies in the management of asthma

